

Crop	Time of application	Dosage
Paddy	At tillering stage	25ml
	20days after first spray	25ml
Wheat	At tillering stage	25ml
	30 to 40 days after 1st spray	25ml
Maize	4 to 10 leaf stage	25ml
	Repeat after 30 days	25ml
Cotton	45 to 60 days after sowing	25ml
	60 to 75 days after first spray	50ml
	75 to 90 days after second spray	50ml
Sugarcane	50 to 60 days after planting	50ml
	30 days after first spray	50ml
Groundnuts	30 days after sowing	25ml
	50days after sowing	25ml
	70days after sowing	25ml
Apples	At flowering	25ml
	At petal fall	25ml
	Two weeks after petal falls	25ml
Grapes	At 3 leaf stage	50ml
	At five leaf stage	50ml
	10 days after first spray	50ml
	Repeat after 12 days	50ml
Pulses	At flowering stage	25ml
	Repeat after 15 days	25ml
Potato	5 to 6 leaf stage	25ml
	15 days after first stage	25ml
	15 days after second spray	50ml
Bhindi	At flowering	25ml
	At fruit setting	50ml

Chilli	20 to 30 days after Transplanting	25ml
	At flowering stage	25ml
	At fruit stage	25ml
Banana	One week before flowering	25ml
Tomato	20 to 25 days after transplanting	25ml
	10 to 15 days after first spray	25ml
	10 to 15 days after first spray	25ml
Peas	Three leaf stage	25ml
	8 to 12 days after first spray	25ml
	8 to 12 days after second spray	25ml
Cabbage & Cauliflower	Five leaf stage	25ml
	15 days after first spray	25ml
	15 days after second spray	25ml
Onion	When plant is 5 – 10 cm high	25ml
	30 days after first spray	25ml
	30 days after second spray	25ml
Ginger & Turmeric	40 days after sowing	25ml
	15 days after first spray	25ml
	15 days after second spray	25ml
Tea	At the time of pruning	25ml
	10 days after first spray	25ml
	10 days after second spray	25ml
Flowers & Ornamental plants	Before Flowering	25ml
Cardamom	At flowering stage	25ml
	At capsule formation	25ml